

**ISSN 2518-1491 (Online),
ISSN 2224-5286 (Print)**

ҚАЗАҚСТАН РЕСПУБЛИКАСЫ
ҰЛТТЫҚ ҒЫЛЫМ АКАДЕМИЯСЫНЫҢ

Д.В. Сокольский атындағы «Жанармай,
катализ және электрохимия институты» АҚ

Х А Б А Р Л А Р Ы

ИЗВЕСТИЯ

НАЦИОНАЛЬНОЙ АКАДЕМИИ НАУК
РЕСПУБЛИКИ КАЗАХСТАН
АО «Институт топлива, катализа и
электрохимии им. Д.В. Сокольского»

NEWS

OF THE ACADEMY OF SCIENCES
OF THE REPUBLIC OF KAZAKHSTAN
JSC «D.V. Sokolsky institute of fuel, catalysis
and electrochemistry»

**SERIES
CHEMISTRY AND TECHNOLOGY**

2 (440)

MARCH – APRIL 2020

PUBLISHED SINCE JANUARY 1947

PUBLISHED 6 TIMES A YEAR

ALMATY, NAS RK

NAS RK is pleased to announce that News of NAS RK. Series of chemistry and technologies scientific journal has been accepted for indexing in the Emerging Sources Citation Index, a new edition of Web of Science. Content in this index is under consideration by Clarivate Analytics to be accepted in the Science Citation Index Expanded, the Social Sciences Citation Index, and the Arts & Humanities Citation Index. The quality and depth of content Web of Science offers to researchers, authors, publishers, and institutions sets it apart from other research databases. The inclusion of News of NAS RK. Series of chemistry and technologies in the Emerging Sources Citation Index demonstrates our dedication to providing the most relevant and influential content of chemical sciences to our community.

Қазақстан Республикасы Ұлттық ғылым академиясы "ҚР ҰҒА Хабарлары. Химия және технология сериясы" ғылыми журналының Web of Science-тің жаңаланған нұсқасы Emerging Sources Citation Index-те индекстелуге қабылданғанын хабарлайды. Бұл индекстелу барысында Clarivate Analytics компаниясы журналды одан әрі the Science Citation Index Expanded, the Social Sciences Citation Index және the Arts & Humanities Citation Index-ке қабылдау мәселесін қарастыруды. Web of Science зерттеушілер, авторлар, баспашилар мен мекемелерге контент тереңдігі мен сапасын ұсынады. ҚР ҰҒА Хабарлары. Химия және технология сериясы Emerging Sources Citation Index-ке енүі біздің қоғамдастық үшін ең өзекті және беделді химиялық ғылымдар бойынша контентке адалдығымызды білдіреді.

НАН РК сообщает, что научный журнал «Известия НАН РК. Серия химии и технологий» был принят для индексирования в Emerging Sources Citation Index, обновленной версии Web of Science. Содержание в этом индексировании находится в стадии рассмотрения компанией Clarivate Analytics для дальнейшего принятия журнала в the Science Citation Index Expanded, the Social Sciences Citation Index и the Arts & Humanities Citation Index. Web of Science предлагает качество и глубину контента для исследователей, авторов, издателей и учреждений. Включение Известия НАН РК в Emerging Sources Citation Index демонстрирует нашу приверженность к наиболее актуальному и влиятельному контенту по химическим наукам для нашего сообщества.

Бас редакторы
х.ғ.д., проф., ҚР ҮҒА академигі
М.Ж. Жұрынов

Редакция алқасы:

Агабеков В.Е. проф., академик (Белорус)
Волков С.В. проф., академик (Украина)
Воротынцев М.А. проф., академик (Ресей)
Газалиев А.М. проф., академик (Қазақстан)
Ергожин Е.Е. проф., академик (Қазақстан)
Жармағамбетова А.К. проф. (Қазақстан), бас ред. орынбасары
Жоробекова Ш.Ж. проф., академик (Қырғыстан)
Иткулова Ш.С. проф. (Қазақстан)
Манташян А.А. проф., академик (Армения)
Пралиев К.Д. проф., академик (Қазақстан)
Баешов А.Б. проф., академик (Қазақстан)
Бұркітбаев М.М. проф., академик (Қазақстан)
Джусипбеков У.Ж. проф., корр.-мүшесі (Қазақстан)
Молдахметов М.З. проф., академик (Қазақстан)
Мансуров З.А. проф. (Қазақстан)
Наурызбаев М.К. проф. (Қазақстан)
Рудик В. проф., академик (Молдова)
Рахимов К.Д. проф., академик (Қазақстан)
Стрельцов Е. проф. (Белорус)
Тәшімов Л.Т. проф., академик (Қазақстан)
Тодераш И. проф., академик (Молдова)
Халиков Д.Х. проф., академик (Тәжікстан)
Фарзалиев В. проф., академик (Әзіrbайжан)

«ҚР ҮҒА Хабарлары. Химия және технология сериясы».

ISSN 2518-1491 (Online),

ISSN 2224-5286 (Print)

Менишіктенуші: «Қазақстан Республикасының Үлттық ғылым академиясы» Республикалық қоғамдық бірлестігі (Алматы қ.).

Қазақстан республикасының Мәдениет пен ақпарат министрлігінің Ақпарат және мұрағат комитетінде 30.04.2010 ж. берілген №1089-Ж мерзімдік басылым тіркеуіне қойылу туралы куәлік.

Мерзімділігі: жылдан 6 рет.

Тиражы: 300 дана.

Редакцияның мекенжайы: 050010, Алматы қ., Шевченко көш., 28; 219, 220 бөл.; тел.: 272-13-19; 272-13-18,
<http://chemistry-technology.kz/index.php/en/arhiv>

© Қазақстан Республикасының Үлттық ғылым академиясы, 2020

Редакцияның мекенжайы: 050100, Алматы қ., Қонаев к-сі, 142, «Д. В. Сокольский атындағы отын, катализ және электрохимия институты» АҚ, каб. 310, тел. 291-62-80, факс 291-57-22, e-mail:orgcat@nursat.kz

Типографияның мекенжайы: «NurNaz GRACE», Алматы қ., Рысқұлов көш., 103.

Г л а в н ы й р е д а к т о р
д.х.н., проф., академик НАН РК
М.Ж. Журинов

Р е д а к ц и о н на я кол л е г и я:

Агабеков В.Е. проф., академик (Беларусь)
Волков С.В. проф., академик (Украина)
Воротынцев М.А. проф., академик (Россия)
Газалиев А.М. проф., академик (Казахстан)
Ергожин Е.Е. проф., академик (Казахстан)
Жармагамбетова А.К. проф. (Казахстан), зам. гл. ред.
Жоробекова Ш.Ж. проф., академик (Кыргызстан)
Иткулова Ш.С. проф. (Казахстан)
Манташян А.А. проф., академик (Армения)
Пралиев К.Д. проф., академик (Казахстан)
Баешов А.Б. проф., академик (Казахстан)
Буркитбаев М.М. проф., академик (Казахстан)
Джусипбеков У.Ж. проф., чл.-корр. (Казахстан)
Мулдахметов М.З. проф., академик (Казахстан)
Мансуров З.А. проф. (Казахстан)
Наурызбаев М.К. проф. (Казахстан)
Рудик В. проф., академик (Молдова)
Рахимов К.Д. проф., академик (Казахстан)
Стрельцов Е. проф. (Беларусь)
Ташимов Л.Т. проф., академик (Казахстан)
Тодераш И. проф., академик (Молдова)
Халиков Д.Х. проф., академик (Таджикистан)
Фарзалиев В. проф., академик (Азербайджан)

«Известия НАН РК. Серия химии и технологии».

[ISSN 2518-1491 \(Online\)](#),

[ISSN 2224-5286 \(Print\)](#)

Собственник: Республиканское общественное объединение «Национальная академия наук Республики Казахстан» (г. Алматы).

Свидетельство о постановке на учет периодического печатного издания в Комитете информации и архивов Министерства культуры и информации Республики Казахстан №10893-Ж, выданное 30.04.2010 г.

Периодичность: 6 раз в год.

Тираж: 300 экземпляров.

Адрес редакции: 050010, г. Алматы, ул. Шевченко, 28; ком. 219, 220; тел. 272-13-19; 272-13-18,
<http://chemistry-technology.kz/index.php/en/arhiv>

© Национальная академия наук Республики Казахстан, 2020

Адрес редакции: 050100, г. Алматы, ул. Кунаева, 142, АО «Институт топлива, катализа и электрохимии им. Д.В. Сокольского», каб. 310, тел. 291-62-80, факс 291-57-22, e-mail:orgcat@nursat.kz

Адрес типографии: «NurNaz GRACE», г. Алматы, ул. Рыскулова, 103.

E d i t o r i n c h i e f
doctor of chemistry, professor, academician of NAS RK
M.Zh. Zhurinov

E d i t o r i a l b o a r d:

Agabekov V.Ye. prof., academician (Belarus)
Volkov S.V. prof., academician (Ukraine)
Vorotyntsev M.A. prof., academician (Russia)
Gazaliyev A.M. prof., academician (Kazakhstan)
Yergozhin Ye.Ye. prof., academician (Kazakhstan)
Zharmagambetova A.K. prof. (Kazakhstan), deputy editor in chief
Zhorobekova Sh.Zh. prof., academician (Kyrgyzstan)
Itkulova Sh.S. prof. (Kazakhstan)
Mantashyan A.A. prof., academician (Armenia)
Praliyev K.D. prof., academician (Kazakhstan)
Bayeshov A.B. prof., academician (Kazakhstan)
Burkitbayev M.M. prof., academician (Kazakhstan)
Dzhusipbekov U.Zh. prof., corr. member (Kazakhstan)
Muldakhmetov M.Z. prof., academician (Kazakhstan)
Mansurov Z.A. prof. (Kazakhstan)
Nauryzbayev M.K. prof. (Kazakhstan)
Rudik V. prof., academician (Moldova)
Rakhimov K.D. prof., academician (Kazakhstan)
Streltsov Ye. prof. (Belarus)
Tashimov L.T. prof., academician (Kazakhstan)
Toderash I. prof., academician (Moldova)
Khalikov D.Kh. prof., academician (Tadzhikistan)
Farzaliyev V. prof., academician (Azerbaijan)

News of the National Academy of Sciences of the Republic of Kazakhstan. Series of chemistry and technology.

ISSN 2518-1491 (Online),

ISSN 2224-5286 (Print)

Owner: RPA "National Academy of Sciences of the Republic of Kazakhstan" (Almaty).

The certificate of registration of a periodic printed publication in the Committee of Information and Archives of the Ministry of Culture and Information of the Republic of Kazakhstan N 10893-Ж, issued 30.04.2010.

Periodicity: 6 times a year.

Circulation: 300 copies.

Editorial address: 28, Shevchenko str., of. 219, 220, Almaty, 050010, tel. 272-13-19; 272-13-18,
<http://chemistry-technology.kz/index.php/en/arhiv>

© National Academy of Sciences of the Republic of Kazakhstan, 2020

Editorial address: JSC «D.V. Sokolsky institute of fuel, catalysis and electrochemistry», 142, Kunayev str., of. 310, Almaty, 050100, tel. 291-62-80, fax 291-57-22, e-mail: orgcat@nursat.kz

Address of printing house: «NurNaz GRACE», 103, Ryskulov str, Almaty.

N E W S

OF THE NATIONAL ACADEMY OF SCIENCES OF THE REPUBLIC OF KAZAKHSTAN

SERIES CHEMISTRY AND TECHNOLOGY

ISSN 2224-5286

<https://doi.org/10.32014/2020.2518-1491.21>

Volume 2, Number 440 (2020), 39 – 46

UDC 547.99

IRSTI 31.23.23

**A.B. Tashimbetova¹, A.K. Umbetova¹, Z.B. Halmenova¹,
Y.S. Ikhsanov¹, M.I. Choudhary², Zh.A. Abilov¹, G.Sh. Burasheva¹**

¹Al-Farabi Kazakh National University Research Institute
of New Technologies and Materials Republic of Kazakhstan, Almaty, Kazakhstan;

²International Center for Chemical and Biological Sciences,
Karachi University, Karachi, Pakistan.

E-mail: erbol.ih@gmail.com, alma_0875@mail.ru, zaure.halmenova@mail.ru, akniyettashimbetova@mail.ru,
abilovs51@mail.ru, iqbal.choudhary@iccs.edu, gauharbur@mail.ru

CHEMICAL STUDY OF *LAVANDULA FERTILIZER*

Abstract. Genus *Lavandula* – specially grown as an aromatic and medicinal plant. In inflorescences, the content of essential oil is from 0.8 % to 2.6 %, in leaves up to 0.3 %. The main components of the essential oil are linalool (10-30 %) in the free state and its esters with acetic, butyric, valeric, caproic acids (30-60 %), as well as geraniol, citral, borneolene, α -pinene and others.

Currently, the plant is used as an ornamental, as a spice in cooking, as well as for medicinal purposes. In traditional medicine, flowers, leaves and branches of lavender are used.

In Bulgaria, lavender is used as a means of calming the nervous system, relaxing baths and for skin diseases.

In Germany, ointments are made from lavender petals. In Austria, lavender leaves are collected before flowering and are used as soothing and anti-inflammatory drugs. In Poland, in combination with a pharmacy chamomile flower, the flowers of a lavender plant are used in the healing of a fresh voice and bronchial disease.

In France, an infusion of lavender flowers is used as an urolithic substance.

The object of the study is raw materials *L.langustifolia* of individual collection at the experimental site of the laboratory of medicinal plants of the Institute of Phyto-Introduction and Botany at the Ministry of Science and Education of the Republic of Kazakhstan of Almaty.

The technology for producing a biologically active complex from the plant species under study has been developed by varying the nature of the extractant, its ratio with raw materials, time and extraction ratio.

The optimal condition for obtaining a biologically active complex from a plant is: extractant – 50 % ethyl alcohol, the ratio of extractant to raw material – 1: 9, the time of double extraction – 72 hours, temperature – 25 °C.

The fractional composition of the aerial mass of *L.langustifolia*, which is represented by water-soluble and water-insoluble fractions, was studied.

The article presents for the first time the data of a qualitative and quantitative analysis of the lipophilic components of the aerial mass of *L.langustifolia*, determined by gas-liquid chromatography with mass spectrometry (GC / MS).

Analyzes were carried out in a laboratory of chemistry of natural coefficients, University of Karachi, Karachi, Pakistan, the structure and quantitative content of 16 compounds were established in the aerial mass of the plant *L.langustifolia*.

The identification of the components was carried out by analogy with the known mass spectra of the samples embedded in the computer data bank and relative retention times. Quantitative determination of the composition of the analyzed mixture was carried out by the method of normalization by peak areas.

Keywords: *Lavandula (L. angustifolia)*, GC-MS, extraction, BAS, aerial parts.

Introduction

Lavandula root is a rod, a tree, and strongly branched on its support. When the new bushes are stretched to the surface, the obsolete bumps bend over. The leaves are rectangular, oval greens, bearded or rectangular, with the edges. The young leaves of the lavanta plant are gray, aged leaves – green [1-3].

Lavandula was specially crafted as aromatic and medicinal plant in the year 2000. Today *Lavandula* plant is used in many different spheres. It is widely used as lavender in lavender, as a spice in culinary, as well for therapeutic purposes, lavender flowers, leaves and branches are used in folk medicine. Collect lavender flowers at the time of flowering and dry it on a sunny day. Then, prepare the flowers, leaves and sprays separately for use in the pots [4-6].

Flowers of *Lavandula* contain uric acid, cinnamic and garnarin. Essential oils will be 0.8-2.6 % in the bouquet, 0.3 % in the leaves and 0.19 % in the class. The main components of the essential oils are linoleum (10-30 %) in vacuum and its esters (30-60 %) with acetic, oily, valerian and caproic acids, as well as alumina, citral, borneol, bisabolene, α -pinene, dihydrochryophyllene epoxy, as healing.

It is known that the composition of plant extracts is directly related to various factors such as the temperature of the extraction process, the extraction time, the plant raw material and the nature of the solvent or solvent system used.

Depending on the tasks to be solved, a wide array of organic solvents such as hexane, chloroform or benzene is used to extract biologically active substances.

However, most organic solvents are highly toxic compounds, have a build-up effect in tissues or are carcinogens, whereby it is required to include additional steps for drug substance purification and additional methods for quality control of such preparations. These measures lead to an imminent increase in the cost of end product [7-11].

This article analyzes the lipophilic composition of the ethanol extract *Lavandula* with various alcohol / water content. The aim of the study was to compare the chemical composition of the extracts from *Lavandula* using two solvents, which are different in polar type, to allow the two systems to be used as a basis for the medicinal forms.

Materials and Methods

The selection of the optimal technological parameters of raw plant materials is aimed at the identification of those affecting directly the composition and the quality of the final complex of biologically active substances (BAS) [12]:

- 1) the nature of the solvents;
- 2) raw material:solvent ratio;
- 3) the temperature;
- 4) the duration of the extraction;
- 5) the extraction frequency.

The selection of the optimal solvent was carried out on the basis of the qualitative analysis of the main BAS groups and the quantitative extraction of extractive substances by pharmacopocial methods [13].

Determination of lipophilic components of a substance by chromatography spectrometry. The extracted extrudate is extracted with hexane (1: 2) for 72 hours at 20-24 °C. The extract is enriched in a rotary evaporator in a soft case.

A Perkin-Elemer Autosystem gas chromatograph with an XL mass-selective detector – TurboMass is used with a flare quartz capillary column (30 x 2.5 mm, 0.25 μ m thick), PE-5 coating and a 99.9 % helium-filled phase. It starts at a temperature of 60 °C (works for 5 minutes), from 20 °C / min to 180 °C, from 3.5 °C/min to 290 °C. The final temperature is maintained for 40 minutes.

Mass spectra are ionized with an emission of 70 eV and a full scan in the range of 40-350 a.m. The injector temperature was 310 °C, and the RA-TiO sample was divided by 1:60 according to the model [14-20].

Methods for determining the qualitative composition of the ground part of plants and roots, determining the composition of fractions, as well as determining the structure and purity of the separation of compounds in raw materials:

- extraction of various polar solvents;

– two-dimensional and one-dimensional chromatography on FN3 paper (Germany) and with a system of solutions.

As the main method for studying chemical composition, the authors opted for the method of high performance gas chromatography with mass-selective detector as the most accurate and universal, which allows for identification of the widest possible spectrum of compounds.

When selecting the optimal extractant, the following solvents are used: ethyl alcohol (50 %, 70 %, 90 %) water solution. 5 g of *Lavandula (L. angustifolia)* of the ground part of the plant, 50 ml capacity, packaged in sausages. The raw material and the solvent obtained in a ratio of 1:6 are kept for 24 hours at room temperature. After 24 hours, the resulting extract can be filtered through a Buchner filter and a vacuum pump. The resulting extract is measured with a measuring cylinder, poured into a porcelain dish, evaporated in a water bath, and the substance is removed.

In the technology of substance separation, the main parameter is the ratio of raw materials and extractant. In order to determine the optimal volume, the selected extractant changes the ratio of raw material and solvent from a ratio of 1:4 to a ratio of 1:10. 5 g crushed (*L. angustifolia*) – ground part of plants of various sizes (20, 30, 40, 45, 50) 50 % extraction with ethyl alcohol. In addition, during the extraction process as a stable factor: extraction time (24 hours) and temperature (20-25 °C).

Determination of technological parameters for obtaining substances


Selecting the extraction state: Obtaining the substance was carried out according to the following scheme:

50-100 g vegetable raw materials *Lavandula (L. angustifolia)* 250-500 ml of 50 % water-ethyl alcohol, preparation of raw materials and extractant in a ratio of 1:9 and settling at room temperature for 72 hours. The contents of the flask are thoroughly mixed, and the paper is filtered into a dry flask through a filter. This extraction process is repeated 2 times using the high-temperature method.

The resulting extracts are added and filtered with paper used at the beginning of the filter.

Results and discussion

According to the results obtained (*L. angustifolia*), during 72 hours of extraction at a temperature of 24-28 °C. the time and multiplicity of extraction are the same. 100 g of dried and standardized plant raw materials were obtained for extraction.



Picture 1 - General provisions of *Lavandula* using various extractants origin of the substance from the ground part of the plant , %

Using the obtained extracts of *Lavandula* (L. from the aboveground part of the angustifolia plant, a substance of 50 % ethyl alcohol was taken in large quantities.

Table 1 - *Lavandula* is related to the relationship between "raw material-extractant" %

Ratio of raw material (g) and extractant (ml)	<i>Lavandula (L.langustifolia)</i> , %
1:4	-
1:6	27,44
1:8	31,96
1:9	55,94
1:10	55,36

Using the selected extractant, the optimal raw material – extractant ratio is 1: 9. if you choose the right ratio of raw materials and extractant, you can save time by knowing the content and ratio of extractant in the industrial industry.

Table 2 - General provisions depending on the time of the extraction process *Lavandula* %

Time of extraction	<i>Lavandula (L.langustifolia)</i> , %
12	27,24
24	55,94
48	75,52
72	88,66

Table 3 - General provisions depending on the multiplicity of extraction of *Lavandula*, %

Number of extractions	<i>Lavandula (L.langustifolia)</i> , %
1 extraction	28,01
2 extraction	28,14

As the next step, the extract was examined by high-performance gas chromatography with mass selective detector Aligent Technologies 6400 Series Triple Quadrupole LC/MS under the following conditions: Poroshell 120 EC-C18 column (50 mm long, 3 mm in diameter, with the coating substance particle size of 4.0, 2.7 and 1.9 μm) was used, with 10 % aqueous solution of methanol as mother solvent and 90 % methanol as final solvent at the pressure of 11.5 mPa and the temperature of 40 °C. Components were identified by mass spectra and retention times using the NIST library and Wiley LC/MS.

Based on the research of the studied plant raw materials, it was found that there are 16 compounds. Plant raw materials contain the amount of methyl ester of linoleic acid (29.34 %) and hexahydrofarrenal acetone (0.41 %). *Lavandula* plant raw material contains methyl ester of linoleic acid (29.34 %), ethyl ether of palmitic acid (21.48 %) and methyl ester of palmitic acid (15.42 %). The *Lavandula* plant contains β -santalol (0.41 %), which contains very small amounts of Sesquiterpenoid compounds.

It is obvious that with in case of an unchanged amount of the plant material, the increase of the extragent amount in the extraction process results in increase of the substance dissolved and transported from the cell to the intercellular space. At the same time, the increase of the volume of the extragent leads to decrease of BAS concentration in the extract.

The ground raw material is introduced to a five-, seven-, nine- and twelve-fold amount of the solvent for studying the dependence of the raw material solvent ratio on the BAS amount in the plants studies. The results are shown in Fig. 1, The raw material solvent ratio has a positive effect. In fact, the solvent in the ratio considered brings about a greater amount of the extractive substances obtained despite the solvent used. This is consistent with the mass transfer principle as the concentration gradient between the solid and most of the liquid is the driving force.

In accordance with the mass transfer law the difference between the concentration of the extract and the solvent increases the transition of the soluble substances into the solvent and continues until the establishment of equilibrium. It is so because the concentration difference is the driving force of the diffusion process.

However, the increase of the solvent consumption is limited by the cost of concentrating the extract obtained.

It follows that the cost of the extract evaporation is an important factor requiring consideration. Therefore, the smaller the raw material: solvent ratio, the more efficient from an economic point of view is the extraction process.

The effect of the extraction temperature on the process of maximum extraction of BAS complex is studied. The temperature increase leads to an increase of the solubility of the various polyphenolic constituents, as well as to an increase of the rate of their diffusion. Since the boiling point of the solvents used does not exceed 80 °C no significant change of the composition of the extracted substances is expected.

Table 4 - Lipophilic composition of *Lavandula angustifolia* plant

No	Connection name	RT (min)	Amount, %	Molecular formula
1	Methyl 4-propan-2-yl benzoate	17.021	0.59	C ₁₁ H ₁₄ O ₂
2	α - terpene	19.403	1.88	C ₁₀ H ₁₆
3	Caryophyllene oxide	20.427	7.87	C ₁₅ H ₂₄ O
4	δ - cedrol	21.173	4.03	C ₁₅ H ₂₆ O
5	Lancel	21.635	0.73	C ₁₅ H ₂₄ O
6	Caryophylline oxide	21.895	2.41	C ₁₅ H ₂₄ O
7	2,6,8- trimethylbicyclo [4.2.0] oct-2-en-1,8-diol	22.02	0.88	C ₁₁ H ₁₈ O ₂
8	β - santalol	22.222	0.7	C ₁₅ H ₂₄ O
9	Hexahydrofarnesyl acetone	24.369	0.41	C ₁₈ H ₃₆ O
10	8-Oxo-2'- deoxyguanosine	25.963	0.63	C ₁₅ H ₂₂ O
11	Palmettic acid methyl ester	26.24	15.42	C ₁₇ H ₃₄ O ₂
12	Palmetinic acid ethyl ester	28.253	21.48	C ₁₈ H ₃₆ O ₂
13	Linoleic acid methyl ester	31.516	29.34	C ₁₉ H ₃₂ O ₂
14	Phytol	31.709	4.25	C ₂₀ H ₄₀
15	Stearic acid methyl ester	31.902	3.98	C ₁₉ H ₃₈ O ₂
16	Linoleic Acid Ethyl Ester	32.506	1.4	C ₂₀ H ₃₄ O ₂

Conclusion

The lipophilic composition of the surface of the *angustifolia Lavandula* plant was determined by gas chromatography using an ion detector and an Agilent HP-5MC device at Karatekin University, Chancara, Turkey. The studied plant material contains linoleic acid methyl ether (29.34 %), palmitic acid ethyl ether (21.48 %) and palmitic acid methyl ether (15.42 %). *Lavandula* contains a very small amount of β-santalol (0.41 %), which is part of sesquiterpenoid.

16 compounds were identified using the chromatogram shown in table 4.(methyl-4-propane-2-yl benzoate, α – terpinene, caryophyllene oxide, δ – cedrol, Lancel, 2,6,8 – trimethylbicyclo [4.2.0]Oct-2-EN-1,8-diol, β – santalol, hexahydrofarnesylacetone, 8-Oxo-2'-deoxyguanosine, methyl palmitic acid ester, ethyl palmitic acid ester, methyl ether linolenic acid, phytol, stearic acid methyl ester, linoleic acid ethyl ester). The resulting lavender essential oil contains 70-80 % saturated C6-c44 hydrocarbons. Quantitative quantities are: docosan (2.8779 %), genacosan (2.4896 %), non-docosan z-14 (2.2156 %), tetracosan (2.3894 %). Butyl-1-methylpropyl ether (4.4067 %) and cyclochloroctane (3.0660 %), 1,2-Benzenesulfonic acid. Test essential oil contains from 10 to 15 esters. In addition, korneol found in essential oils (0,0287 %) and caryophyllia (0,0123 %).

**А.Б. Ташимбетова¹, А.К. Умбетова¹, З.Б. Халменова¹,
Е.С. Ихсанов, М.И. Чоудхари², Ж.А. Абилов¹, Г.Ш. Бурашева¹**

¹ әл-Фараби атындағы Қазақ ұлттық университеті,
Жаңа технологиялар мен материалдар ғылыми-зерттеу институты
Қазақстан Республикасы, Алматы, Қазақстан;

² Халықаралық химиялық және биологиялық ғылымдар орталығы,
Карачи университеті, Карачи, Пәкістан

LAVANDULA FERTILIZER-ДІҢ ХИМИЯЛЫҚ ЗЕРТТЕУІ

Аннотация. Лаванда (*Lavandula*) біздің дәуірімізге дейін хош иісті және дәрілік өсімдік ретінде арнағы өсірілген. Эфир майы құрамы гүл шоғырында – 0,8 %-дан 2,6 %-га дейін, жапырактарында 0,3 %-га дейін болады. Эфир майының негізгі компоненттерінде еркін жағдайдағы линалоол (10-30 %) және сірке сусы, май, валериян, капрон қышқылы (30-60 %) бар, сондай-ақ оның эфирлері – гераниол, цитраль, борнеолен, бисаболен, α-пинен. Қазіргі уақытта өсімдік өнімдерде, дәмдеуіш ретінде аспаздық үшін, сондай-ақ емдік үшін де пайдаланылады. Халықтық медицинада дәрілік максаттармен лаванданың ғулдері, жапырактары мен бұтақтары қолданылады, бұтақтарын бөлек бөтелкелерге жинап, пайдалануға дайындауды. Болгарияда лаванда жүйке жүйесін тыныштандыратын, ванналар мен тері ауруларын емдейтін дәрі ретінде қолданылады. Германияда лаванда жапырактарынан жақпа жасайды. Австрияда лаванда жапырактары гүлденгенге дейін жиналады, олар тыныштандыратын және қабынуға карсы препараттар ретінде пайдаланылады. Польшада дәріханалық түймедақ ғулімен бірге, лаванда өсімдіктерінің ғулдері жаңа дауыс пен бронх ауруын емдеуде қолданылады. Францияда лаванда ғүлінің тұнбасы несеп айдайтын зат ретінде пайдаланылады.

Зерттеу объектісі – Қазақстан Республикасы Ғылым және Білім Министрлігі жаңындағы фитоинтродукция және ботаника институты дәрілік өсімдіктер зертханасының эксперименталдық бөлімінде жеке жиналған *Lavandula (L.langustifolia)* шикізаты.

Зерттеліп жатқан өсімдік түрлерінен биологиялық белсенді кешен алу технологиясы экстракция затының табигатын, шикізатқа қатынасын, уақытты және экстракция коэффициентін өзгерте отырып жасалды.

Өсімдіктен биологиялық белсенді кешен алудың онтайлы шарты: экстрагент – 50 % этил спирті, экстракцияның шикізатқа қатынасы – 1: 9, қосарланған экстракция уақыты – 72 сағат, температура – 250 °C.

L.langustifolia жер үсті массасының фракциялық құрамы зерттелді, ол суда еритін және суда ерімейтін фракциялардан тұрады.

Мақалада алғаш рет *L.langustifolia* жер үсті массасының липофильді компоненттерінің сапалық және сандық талдауының деректері келтірілді, масс-спектрометриясы (GC/MS) бар газ сүйіктықты хроматография әдісімен анықталған.

Талдаулар табиги қосылыстар химиясы зертханасында, Карачи қаласының университетінде, Пәкістан қаласында *L.langustifolia* өсімдігінің жер үсті массасында 16 қосылыстың құрылымы мен сандық құрамы орнатылды.

Компоненттерді сәйкестендіру компьютердің деректер банкіне салынған үлгілердің белгілі масс-спектрлеріне және ұстап қалу уақытына ұқсас жүзеге асырылады. Талданатын қоспаның құрамын сандық анықтау шындар ауданы бойынша қалыпқа келтіру әдісімен жүзеге асырылды.

**А.Б. Ташимбетова¹, А.К. Умбетова¹, З.Б. Халменова¹,
Е.С. Ихсанов, М.И. Чоудхари², Ж.А. Абилов¹, Г.Ш. Бурашева¹**

¹ Казахский национальный университет имени аль-Фараби
Научно-исследовательский институт новых технологий и материалов
Республика Казахстан, Алматы, Казахстан;

²Международный центр химических и биологических наук университета Карачи,
г. Карачи, Пакистан

ХИМИЧЕСКОЕ ИССЛЕДОВАНИЕ LAVANDULA FERTILIZER

Аннотация. Род Лаванда (*Lavandula*) – специально выращенный как ароматное и лекарственное растение. В соцветиях содержание эфирного масла от 0,8 % до 2,6%, в листьях до 0,3 %. Основными компонентами эфирного масла являются линалоол (10–30%) в свободном состоянии и его эфиры с уксусной,

масляной, валерияновой, капроновой кислотами (30–60 %), а также гераниол, цитраль, борнеолен, бисаболен, α -пинен и другие.

В настоящее время растение используется как декоративное, в качестве специй в кулинарии, а также в лечебных целях. В народной медицине используются цветы, листья и ветви лаванды.

В Болгарии лаванду используют как средство, успокаивающее нервную систему, расслабляющие ванны и при кожных заболеваниях.

В Германии делают мази из лепестков лаванды. В Австрии собраны листья лаванды до цветения и используются в качестве успокаивающих и противовоспалительных препаратов. В Польше, в сочетании с аптечным цветком ромашки, цветки лавандового растения применяются при заживлении свежей раны и бронхиальной болезни.

Во Франции настой цветков лаванды используют как мочегонное вещество.

Объектом исследования является сырье *Lavandula (L.langustifolia)* индивидуального сбора на экспериментальном участке лаборатории лекарственных растений Института фитоинтродукции и ботаники при Министерстве науки и образования Республики Казахстан города Алматы.

Отработана технология получения биологически активного комплекса из исследуемого вида растения варированием природы экстрагента, его соотношением с сырьем, времени и кратности экстракции.

Оптимальным условием для получения биологически активного комплекса из растения является: экстрагент – 50 % этиловый спирт, соотношение экстрагента и сырья – 1:9, время двухкратной экстракции – 72 часа, температура – 25°C.

Изучен фракционный состав надземной массы *L.langustifolia*, который представлен водорастворимыми и нерастворимыми в воде фракциями.

В статье впервые приведены данные качественного и количественного анализа липофильных компонентов надземной массы *L.langustifolia*, определены методом газожидкостной хроматографии с масс-спектрометрией (GC/MS).

Анализы проведены в лаборатории химии природных соединений, Университета города Карачи, г. Карачи, Пакистан, в надземной массе растения *L.langustifolia* установлены структуры и количественное содержание 16 соединений.

Идентификация компонентов осуществлена по аналогии с известными масс-спектрами образцов, заложенных в банк данных компьютера и относительным временем удерживания. Количественное определение состава анализируемой смеси осуществлено методом нормализации по площадям пиков.

Ключевые слова: *Lavandula (L. angustifolia)*, ГХ-МС, экстракция, БАВ, надземная часть.

Information about authors:

Ikhsanov Yerbol Saginovich - PhD Department of Chemistry and Chemical Technology of the Al-Farabi Kazakh National University Kazakh National University, e-mail: erbol.ih@gmail.com, <https://orcid.org/0000-0003-4640-9584>;

Umbetova Almagul Kengebaeva - Candidate of chemical Sciences, faculty of chemistry and chemical technology, al-Farabi Kazakh national University, El. mail: alma_0875@mail.ru, <https://orcid.org/0000-0001-9879-5398>;

Khalmenova Zaure Bisentaeva - Candidate of technical Sciences, faculty of chemistry and chemical technology, al-Farabi Kazakh national University, El. mail: zaure.halmenova@mail.ru, <https://orcid.org/0000-0002-6991-560X>;

Tashimbetova Akniyet Bakytbekkyzy - faculty of chemistry and chemical technology, al-Farabi Kazakh national University, Эл. почта: akniyettashimbetova@mail.ru, <https://orcid.org/0000-0002-0010-805X>;

Abilov Zharkasyn Abduahitovich - Doctor of Chemical Sciences, Professor of the Department of Chemistry and Chemical Technology of Al-Farabi Kazakh National University, Email mail: abilovs51@mail.ru, <https://orcid.org/0000-0002-2665-2539>

Muhammad Ihbal Chowdhari - Doctor of Chemical Sciences, Professor, Karachi University, Email mail: iqbal.choudhary@iccs.edu, <https://orcid.org/0000-0001-5356-3585>;

Burasheva Gauhar Shahmanovna - Doctor of Science professor, Department of Chemistry and Chemical Technology of the Al-Farabi Kazakh National University, e-mail: gauharbur@mail.ru, <https://orcid.org/0000-0003-2935-3531>

REFERENCES

- [1] Pavlov N.B. (1961) Flora Of Kazakhstan. 5th ed. Alma-ATA: an KazSSR. 70-72 p. 3. Beklemishev N.D. Drugs from plants. Almaty, 2002. 208 p.
- [2] Bajtenov M.S. (2001) Flora Kazahstana. Rodovoj kompleks flory. Almaty: Gylym. ISBN 9965-07-036-9
- [3] Mahmedova.M.(1991) Labiate Central Asia. Tashkent. 54-55.
- [4] Uzakbay S.A., Khalmenova V.D., Umbetova A.K. (2016) Investigation of chemical composition of plants *Ogidapim vulgare* in Almaty region. Izvestiya NAS RK. No. 6 (420). P 99-102.
- [5] Ishmuratova M., Tleukenov S.U. (2007) To the study of plants of the family Labiate of the flora of Central Kazakhstan // Proceedings of int.science. Conf. The plant world and its protection. Almaty. P. 352-354.
- [6] Kurgan S.A. (1995) Useful herbs and rare flowers in the garden. Moscow: Nauka. p.128.
- [7] Wulf E.V., Pigulevsky G.V., Albrecht E.A.(1999) Culture of plants giving essential oils on the southern coast of Crimea. Yalta. 126 p.

- [8] Voronina E. P.(1993) introduction of lavender In the main Botanical garden of the Russian Academy of Sciences.GL. Botan.garden's. Vol.168. P. 3-8.
- [9] Romanenko L. G.(1992) Improvement and development of methods of selection of narrow-leaved lavender (*Lavandula angustifolia* Mill.). Auto-Ref-Doct.SEL.sciences'. Moscow, 44 p.
- [10] Buyukli M.V.(1990) Lavender and its culture in the USSR. Chisinau: Carta Moldoveneasca p-327.
- [11] Karasavidis A.O. (2005) Some types of odoriferous raw material in medical practice// Vestnik of VSU.No. 1. p. 205-211.
- [12] Trellakis S, et al.(2012) Subconscious olfactory influences of stimulant and relaxant odors on immune function. Eur Arch Otorhinolaryngol.
- [13] Kim, N.-S., & Lee, D.-S. (2002). Comparison of different extraction methods for the analysis of fragrances from *Lavandula* species by gas chromatography–mass spectrometry. Journal of Chromatography A, 982(1), 31-47. doi:10.1016/s0021-9673(02)01445-0.
- [14] Lu Huil,2, Li He 2, Lu Huan 1, Li XiaoLan3* and Zhou Ai Guo (2010) Chemical composition of laverder essential oil and its andioxidant activity and inhibition against rhinitisrelated bacteria. African Journal of microbiology. Vol. 4. №4. P. 309-313.
- [15] Tabopda, T., Ngoupayo, J., Khan Tanoli, S., Mitaine-Offer, A.-C., Ngadjui, B., Ali, M., Lacaille-Dubois, M.-A. (2009). Antimicrobial Pentacyclic Triterpenoids from *Terminalia superba*. *Planta Medica*, 75(05), 522–527. doi:10.1055/s-0029-1185328
- [16] Beens J., Brinkman U.A.T. Analyst (Cambridge, UK), 2005. Vol. 130. P. 123.
- [17] Aitkulova RE, Abubakirova AA, Kudasova DE, Kaldybekova GM (2016) Role of medicinal plants from south-kazakhstan region for addition into livestock's fodder. News of the national academy of sciences of the republic of kazakhstan series of biological and medical 314: 155–158. (in Eng). <https://doi.org/10.32014/2018.2518-1629>
- [18] Tursynova A., Sunnenova N.S., Erezhepova N, Sarsenbayeva N.B., Kalekeshov A.M., Makashev E.K. (2016) Effect of the feed additive based on bentonite and chlorella body of agricultural animals and birds, News of the national academy of sciences of the republic of Kazakhstan series of biological and medical. 3: 27–33. <https://doi.org/10.32014/2018.2224-526> (in Eng).
- [19] Oscanov B.S., Iksanov Y.S. , Litvinenko Yu.A., Adekenov S.M., Burasheva G.Sh., Biologically active substances from plant sudaeda vera and their anestizing activity News of the national academy of sciences of the republic of kazakhstan series of biological and medical ISSN 2224-5308. Volume 5, Number 329 (2018), 63 – 66. <https://doi.org/10.32014/2018.2518-1629.8>
- [20] Aitkulova R.E., Abubakirova A.A., Kudasova D.E., Kaldybekova G.M. (2016) Role of medicinal plants from south-kazakhstan region for addition into livestock's fodder. News of the national academy of sciences of the republic of kazakhstan series of biological and medical 314: 155 – 158. (in Eng). <https://doi.org/10.32014/2018.2518-1629>

Publication Ethics and Publication Malpractice in the journals of the National Academy of Sciences of the Republic of Kazakhstan

For information on Ethics in publishing and Ethical guidelines for journal publication see <http://www.elsevier.com/publishingethics> and <http://www.elsevier.com/journal-authors/ethics>.

Submission of an article to the National Academy of Sciences of the Republic of Kazakhstan implies that the described work has not been published previously (except in the form of an abstract or as part of a published lecture or academic thesis or as an electronic preprint, see <http://www.elsevier.com/postingpolicy>), that it is not under consideration for publication elsewhere, that its publication is approved by all authors and tacitly or explicitly by the responsible authorities where the work was carried out, and that, if accepted, it will not be published elsewhere in the same form, in English or in any other language, including electronically without the written consent of the copyright-holder. In particular, translations into English of papers already published in another language are not accepted.

No other forms of scientific misconduct are allowed, such as plagiarism, falsification, fraudulent data, incorrect interpretation of other works, incorrect citations, etc. The National Academy of Sciences of the Republic of Kazakhstan follows the Code of Conduct of the Committee on Publication Ethics (COPE), and follows the COPE Flowcharts for Resolving Cases of Suspected Misconduct (http://publicationethics.org/files/u2/New_Code.pdf). To verify originality, your article may be checked by the Cross Check originality detection service <http://www.elsevier.com/editors/plagdetect>.

The authors are obliged to participate in peer review process and be ready to provide corrections, clarifications, retractions and apologies when needed. All authors of a paper should have significantly contributed to the research.

The reviewers should provide objective judgments and should point out relevant published works which are not yet cited. Reviewed articles should be treated confidentially. The reviewers will be chosen in such a way that there is no conflict of interests with respect to the research, the authors and/or the research funders.

The editors have complete responsibility and authority to reject or accept a paper, and they will only accept a paper when reasonably certain. They will preserve anonymity of reviewers and promote publication of corrections, clarifications, retractions and apologies when needed. The acceptance of a paper automatically implies the copyright transfer to the National Academy of Sciences of the Republic of Kazakhstan.

The Editorial Board of the National Academy of Sciences of the Republic of Kazakhstan will monitor and safeguard publishing ethics.

Правила оформления статьи для публикации
в журнале смотреть на сайте:

www:nauka-nanrk.kz

<http://chemistry-technology.kz/index.php/en/arhiv>

ISSN 2518-1491 (Online), ISSN 2224-5286 (Print)

Редакторы: *М. С. Ахметова, Г. Б. Халидуллаева, Д. С. Аленов*
Верстка на компьютере *А.М. Кульгинбаевой*

Подписано в печать 29.03.2020.
Формат 60x881/8. Бумага офсетная. Печать – ризограф.
10 п.л. Тираж 300. Заказ 2.

Национальная академия наук РК
050010, Алматы, ул. Шевченко, 28, т. 272-13-18, 272-13-19