

**ISSN 2518-1491 (Online),
ISSN 2224-5286 (Print)**

ҚАЗАҚСТАН РЕСПУБЛИКАСЫ
ҰЛТТЫҚ ҒЫЛЫМ АКАДЕМИЯСЫНЫҢ

Д.В. Сокольский атындағы «Жанармай,
катализ және электрохимия институты» АҚ

Х А Б А Р Л А Р Ы

ИЗВЕСТИЯ

НАЦИОНАЛЬНОЙ АКАДЕМИИ НАУК
РЕСПУБЛИКИ КАЗАХСТАН
АО «Институт топлива, катализа и
электрохимии им. Д.В. Сокольского»

NEWS

OF THE ACADEMY OF SCIENCES
OF THE REPUBLIC OF KAZAKHSTAN
JSC «D.V. Sokolsky institute of fuel, catalysis
and electrochemistry»

**SERIES
CHEMISTRY AND TECHNOLOGY**

2 (440)

MARCH – APRIL 2020

PUBLISHED SINCE JANUARY 1947

PUBLISHED 6 TIMES A YEAR

ALMATY, NAS RK

NAS RK is pleased to announce that News of NAS RK. Series of chemistry and technologies scientific journal has been accepted for indexing in the Emerging Sources Citation Index, a new edition of Web of Science. Content in this index is under consideration by Clarivate Analytics to be accepted in the Science Citation Index Expanded, the Social Sciences Citation Index, and the Arts & Humanities Citation Index. The quality and depth of content Web of Science offers to researchers, authors, publishers, and institutions sets it apart from other research databases. The inclusion of News of NAS RK. Series of chemistry and technologies in the Emerging Sources Citation Index demonstrates our dedication to providing the most relevant and influential content of chemical sciences to our community.

Қазақстан Республикасы Ұлттық ғылым академиясы "ҚР ҰҒА Хабарлары. Химия және технология сериясы" ғылыми журналының Web of Science-тің жаңаланған нұсқасы Emerging Sources Citation Index-те индекстелуге қабылданғанын хабарлайды. Бұл индекстелу барысында Clarivate Analytics компаниясы журналды одан әрі the Science Citation Index Expanded, the Social Sciences Citation Index және the Arts & Humanities Citation Index-ке қабылдау мәселесін қарастыруды. Web of Science зерттеушілер, авторлар, баспашилар мен мекемелерге контент тереңдігі мен сапасын ұсынады. ҚР ҰҒА Хабарлары. Химия және технология сериясы Emerging Sources Citation Index-ке енүі біздің қоғамдастық үшін ең өзекті және беделді химиялық ғылымдар бойынша контентке адалдығымызды білдіреді.

НАН РК сообщает, что научный журнал «Известия НАН РК. Серия химии и технологий» был принят для индексирования в Emerging Sources Citation Index, обновленной версии Web of Science. Содержание в этом индексировании находится в стадии рассмотрения компанией Clarivate Analytics для дальнейшего принятия журнала в the Science Citation Index Expanded, the Social Sciences Citation Index и the Arts & Humanities Citation Index. Web of Science предлагает качество и глубину контента для исследователей, авторов, издателей и учреждений. Включение Известия НАН РК в Emerging Sources Citation Index демонстрирует нашу приверженность к наиболее актуальному и влиятельному контенту по химическим наукам для нашего сообщества.

Бас редакторы
х.ғ.д., проф., ҚР ҮҒА академигі
М.Ж. Жұрынов

Редакция алқасы:

Агабеков В.Е. проф., академик (Белорус)
Волков С.В. проф., академик (Украина)
Воротынцев М.А. проф., академик (Ресей)
Газалиев А.М. проф., академик (Қазақстан)
Ергожин Е.Е. проф., академик (Қазақстан)
Жармағамбетова А.К. проф. (Қазақстан), бас ред. орынбасары
Жоробекова Ш.Ж. проф., академик (Қырғыстан)
Иткулова Ш.С. проф. (Қазақстан)
Манташян А.А. проф., академик (Армения)
Пралиев К.Д. проф., академик (Қазақстан)
Баешов А.Б. проф., академик (Қазақстан)
Бұркітбаев М.М. проф., академик (Қазақстан)
Джусипбеков У.Ж. проф., корр.-мүшесі (Қазақстан)
Молдахметов М.З. проф., академик (Қазақстан)
Мансуров З.А. проф. (Қазақстан)
Наурызбаев М.К. проф. (Қазақстан)
Рудик В. проф., академик (Молдова)
Рахимов К.Д. проф., академик (Қазақстан)
Стрельцов Е. проф. (Белорус)
Тәшімов Л.Т. проф., академик (Қазақстан)
Тодераш И. проф., академик (Молдова)
Халиков Д.Х. проф., академик (Тәжікстан)
Фарзалиев В. проф., академик (Әзірбайжан)

«ҚР ҮҒА Хабарлары. Химия және технология сериясы».

ISSN 2518-1491 (Online),

ISSN 2224-5286 (Print)

Менишіктенуші: «Қазақстан Республикасының Үлттық ғылым академиясы» Республикалық қоғамдық бірлестігі (Алматы қ.).

Қазақстан республикасының Мәдениет пен ақпарат министрлігінің Ақпарат және мұрағат комитетінде 30.04.2010 ж. берілген №1089-Ж мерзімдік басылым тіркеуіне қойылу туралы куәлік.

Мерзімділігі: жылдан 6 рет.

Тиражы: 300 дана.

Редакцияның мекенжайы: 050010, Алматы қ., Шевченко көш., 28; 219, 220 бөл.; тел.: 272-13-19; 272-13-18,
<http://chemistry-technology.kz/index.php/en/arhiv>

© Қазақстан Республикасының Үлттық ғылым академиясы, 2020

Редакцияның мекенжайы: 050100, Алматы қ., Қонаев к-сі, 142, «Д. В. Сокольский атындағы отын, катализ және электрохимия институты» АҚ, каб. 310, тел. 291-62-80, факс 291-57-22, e-mail:orgcat@nursat.kz

Типографияның мекенжайы: «NurNaz GRACE», Алматы қ., Рысқұлов көш., 103.

Г л а в н ы й р е д а к т о р
д.х.н., проф., академик НАН РК
М.Ж. Журинов

Р е д а к ц и о н на я кол л е г и я:

Агабеков В.Е. проф., академик (Беларусь)
Волков С.В. проф., академик (Украина)
Воротынцев М.А. проф., академик (Россия)
Газалиев А.М. проф., академик (Казахстан)
Ергожин Е.Е. проф., академик (Казахстан)
Жармагамбетова А.К. проф. (Казахстан), зам. гл. ред.
Жоробекова Ш.Ж. проф., академик (Кыргызстан)
Иткулова Ш.С. проф. (Казахстан)
Манташян А.А. проф., академик (Армения)
Пралиев К.Д. проф., академик (Казахстан)
Баешов А.Б. проф., академик (Казахстан)
Буркитбаев М.М. проф., академик (Казахстан)
Джусипбеков У.Ж. проф., чл.-корр. (Казахстан)
Мулдахметов М.З. проф., академик (Казахстан)
Мансуров З.А. проф. (Казахстан)
Наурызбаев М.К. проф. (Казахстан)
Рудик В. проф., академик (Молдова)
Рахимов К.Д. проф., академик (Казахстан)
Стрельцов Е. проф. (Беларусь)
Ташимов Л.Т. проф., академик (Казахстан)
Тодераш И. проф., академик (Молдова)
Халиков Д.Х. проф., академик (Таджикистан)
Фарзалиев В. проф., академик (Азербайджан)

«Известия НАН РК. Серия химии и технологии».

[ISSN 2518-1491 \(Online\)](#),

[ISSN 2224-5286 \(Print\)](#)

Собственник: Республиканское общественное объединение «Национальная академия наук Республики Казахстан» (г. Алматы).

Свидетельство о постановке на учет периодического печатного издания в Комитете информации и архивов Министерства культуры и информации Республики Казахстан №10893-Ж, выданное 30.04.2010 г.

Периодичность: 6 раз в год.

Тираж: 300 экземпляров.

Адрес редакции: 050010, г. Алматы, ул. Шевченко, 28; ком. 219, 220; тел. 272-13-19; 272-13-18,
<http://chemistry-technology.kz/index.php/en/arhiv>

© Национальная академия наук Республики Казахстан, 2020

Адрес редакции: 050100, г. Алматы, ул. Кунаева, 142, АО «Институт топлива, катализа и электрохимии им. Д.В. Сокольского», каб. 310, тел. 291-62-80, факс 291-57-22, e-mail:orgcat@nursat.kz

Адрес типографии: «NurNaz GRACE», г. Алматы, ул. Рыскулова, 103.

E d i t o r i n c h i e f
doctor of chemistry, professor, academician of NAS RK
M.Zh. Zhurinov

E d i t o r i a l b o a r d:

Agabekov V.Ye. prof., academician (Belarus)
Volkov S.V. prof., academician (Ukraine)
Vorotyntsev M.A. prof., academician (Russia)
Gazaliyev A.M. prof., academician (Kazakhstan)
Yergozhin Ye.Ye. prof., academician (Kazakhstan)
Zharmagambetova A.K. prof. (Kazakhstan), deputy editor in chief
Zhorobekova Sh.Zh. prof., academician (Kyrgyzstan)
Itkulova Sh.S. prof. (Kazakhstan)
Mantashyan A.A. prof., academician (Armenia)
Praliyev K.D. prof., academician (Kazakhstan)
Bayeshov A.B. prof., academician (Kazakhstan)
Burkitbayev M.M. prof., academician (Kazakhstan)
Dzhusipbekov U.Zh. prof., corr. member (Kazakhstan)
Muldakhmetov M.Z. prof., academician (Kazakhstan)
Mansurov Z.A. prof. (Kazakhstan)
Nauryzbayev M.K. prof. (Kazakhstan)
Rudik V. prof., academician (Moldova)
Rakhimov K.D. prof., academician (Kazakhstan)
Streltsov Ye. prof. (Belarus)
Tashimov L.T. prof., academician (Kazakhstan)
Toderash I. prof., academician (Moldova)
Khalikov D.Kh. prof., academician (Tadzhikistan)
Farzaliyev V. prof., academician (Azerbaijan)

News of the National Academy of Sciences of the Republic of Kazakhstan. Series of chemistry and technology.

ISSN 2518-1491 (Online),

ISSN 2224-5286 (Print)

Owner: RPA "National Academy of Sciences of the Republic of Kazakhstan" (Almaty).

The certificate of registration of a periodic printed publication in the Committee of Information and Archives of the Ministry of Culture and Information of the Republic of Kazakhstan N 10893-Ж, issued 30.04.2010.

Periodicity: 6 times a year.

Circulation: 300 copies.

Editorial address: 28, Shevchenko str., of. 219, 220, Almaty, 050010, tel. 272-13-19; 272-13-18,
<http://chemistry-technology.kz/index.php/en/arhiv>

© National Academy of Sciences of the Republic of Kazakhstan, 2020

Editorial address: JSC «D.V. Sokolsky institute of fuel, catalysis and electrochemistry», 142, Kunayev str., of. 310, Almaty, 050100, tel. 291-62-80, fax 291-57-22, e-mail: orgcat@nursat.kz

Address of printing house: «NurNaz GRACE», 103, Ryskulov str, Almaty.

N E W S

OF THE NATIONAL ACADEMY OF SCIENCES OF THE REPUBLIC OF KAZAKHSTAN

SERIES CHEMISTRY AND TECHNOLOGY

ISSN 2224-5286

<https://doi.org/10.32014/2020.2518-1491.25>

Volume 2, Number 440 (2020), 69 – 74

UDC 547.99

IRSTI 31.23.23

Y.S. Iksanov, G.E. Tasmagambetova, Y.A. Litvinenko, G.Sh. Burasheva, G.A. Seitimova

Al-Farabi Kazakh National University

Center of Physico-chemical Methods of Research and Analysis

Republic of Kazakhstan, Almaty, Kazakhstan

E-mail: erbol.ih@gmail.com, rumex1978@gmail.com, sitigulnaz@mail.ru,
gauharbur@mail.ru, guljaina_98.14@mail.ru**PHYTOCHEMICAL COMPOSITION OF LIPOPHILIC FRACTION
OF PLANTS OF THE PLANT ROSA CANINA L. GENUS ROSA**

Abstract. The genus of wild rose, or *Rosa L.*, belongs to the family Rosaceae. It has many cultural forms, bred under the general name *Rosa*. At the moment, more than 400 species of rose hips are known. Usually these are upright shrubs, less often creepers, sometimes low tree-like forms or almost herbaceous plants. Rosehip is common in the temperate and subtropical zones of the Northern Hemisphere, it can often be found in the mountainous regions of the tropical zone. Some species of rose hips are common from the Arctic Circle in the north to Ethiopia in the south. On the American continent - from Canada to Mexico, Favorable conditions for Rosehip are in the Mediterranean region. Several species of the Rosehip genus have an extensive distribution area.

Kazakhstan has significant resources of medicinal raw materials of plant origin. Among the most priority representatives of the wild flora are species of the rosehip genus - *Rosa L.* as a most valuable medicinal plant. In total, 21 species of wild rose grow in the republic, including 5 in the central Kazakhstan: *R. glabrifolia* - sh. naked, *R. laxa* Retz. - w. loose, *R. acicularis* Lindl. - (*W. needlefish*), *R. majalis* Herrm. (*R. cinnamomea* L.) - sh. May (*W. cinnamon*) and *R. pimpinellifolia* L. (*R. spinosissima* L.) - W. femoral cell. Kazakhstan species of the genus *Rosa L.*

Rose hips are characterized by a high content of biologically active substances and are widely used both in medicine and in cooking.

In this paper, a comparative analysis of rose hips, *Rosa canina L.*, harvested in the fall of 2018 is carried out.

Plant raw materials are collected in the southern region of the Republic of Kazakhstan (Almaty region). The crushed air-dry raw materials were extracted with hexane and chloroform in the ratio of raw materials-reagent (1:10) in a Soxhlet apparatus. The resulting extract was concentrated under mild conditions to a thick concentrate, which was analyzed on a gas chromatograph with a mass selective detector.

The analyzes were carried out on a gas chromatograph with a mass spectrometric detector 6890N / 5973C (Agilent, USA). The peaks detected in the chromatograms were identified using the NIST'11 and Wiley 10 mass spectral libraries.

The article first studied and conducted a comparative analysis of lipophilic substances contained in the fruits of the plant *Rosa canina L.*, harvested in the Almaty region.

The extracts mainly include hydrocarbons, derivatives of higher carboxylic acids, higher alcohols and some other substances, in particular butyl tetradecl ether of hydrochloric acid, [4- (2-methylpentanoylsulfamoyl) phenyl] amide of 2-methylpentanoic acid in hexane extract and, butyl undecyl ether sulfuric acid, eicosyl trifluoroacetate and ethoxycarbonyl-3-methyl-4-azafluorenone, 2-fluorenilim in the chloroform extract.

Further work will be continued in order to determine the potential biological activity of the obtained extracts.

Keywords: *Rosa canina L.*, СКФ-экстракция, никотин, GC-MS, химический состав.

Introduction

The rosehip genus, *Rosa L.*, belongs to the family Rosaceae. It has many cultural forms, bred under the general name *Rosa*. At the moment, more than 400 species of rose hips are known. Usually these are upright shrubs, less often creepers, sometimes low tree-like forms or almost herbaceous plants. Rosehip is common in the temperate and subtropical zones of the Northern Hemisphere, it can often be found in the mountainous regions of the tropical zone. Some species of rose hips are common from the Arctic Circle in the north to Ethiopia in the south. On the American continent - from Canada to Mexico, Favorable conditions for Rosehip are in the Mediterranean region. Several species of the Rosehip genus have an extensive distribution area.

the north to Ethiopia in the south. On the American continent - from Canada to Mexico. Favorable conditions for Rosehip are in the Mediterranean region. Several species of the Rosehip genus have an extensive distribution area [1-3].

Kazakhstan has significant resources of medicinal raw materials of plant origin. Among the most priority representatives of the wild flora are species of the rosehip genus - *Rosa L.* as a most valuable medicinal plant. In total, 21 species of wild rose grow in the republic, including 5 in the central Kazakhstan: *R. glabrifolia* - sh. naked, *R. laxa Retz.* - w. loose, *R. acicularis Lindl.* - (W. needlefish), *R. majalis Herrm.* (*R. cinnamomea L.*) - sh. May (W. cinnamon) and *R. pimpinellifolia L.* (*R. spinosissima L.*) - W. femoral cell. Kazakhstan species of the genus *Rosa L* [4-5].

Rosehip (canine) (*R. canina*). Branched tall shrub with curved, sparse spikes with single flowers of pale pink or white color. Fruits are smooth, fleshy, light or bright red, spherical or oval-spherical elongated-oval; the flesh is sweet, edible, thick. Sepals fall off before ripening. The pulp content in relation to the weight of the dry fruit is 54.9-65%; seed content 35.0-45.1%. The fruits of this type of rosehip are a weak source of vitamin C. The content of ascorbic acid in the pulp of fruits is usually extremely limited - from 0 to 0.95% and carotene 3.8-12.9 mg% [6].

In this article, we consider lipophilic substances in the composition of *R. canina* fruits identified in hexane and chloroform extracts.

Materials and Methods

Plant raw materials are collected in the southern region of the Republic of Kazakhstan (Almaty region). The crushed air-dry raw materials (500 g) were extracted with hexane in the ratio of raw materials-reagent (1:10) in a Soxhlet apparatus. The extract obtained was concentrated under mild conditions (water bath temperature 40-45 °C) using a vacuum water-jet pump, to a thick concentrate, which was analyzed on a gas chromatograph with a mass-selective detector. The analyzes were performed on a gas chromatograph with a 6890N / 5973C mass spectrometric detector (Agilent, USA) equipped with a Combi-PAL autosampler (CTC Analytics, Switzerland). For GC-MS analysis, 1.00 µl of the sample was injected into the injector of a gas chromatograph using an autosampler at an injector temperature of 250°C. Chromatography was performed using an HP-5ms capillary column (Agilent, USA) with a length of 30 m, an inner diameter of 0.25 mm, and a film thickness of 0.25 µm at a constant carrier gas velocity (helium,> 99.995%, Orenburg-Tekhgaz, Russia) equal to 1.0 ml / min. The program for heating the chromatographic column: holding for 5 min at 40 °C, heating at a speed of 10 °C / min to 280 °C, holding for 5 min. The total chromatographic time was 34 minutes. The temperatures of the quadrupole and the MSD ion source were 150 and 230 °C, respectively [7-16].

Mass spectrometric detection was carried out in the ion scanning mode in the m / z range from 40 to 550 with a solvent delay of 5 min. The peaks detected in the chromatograms were identified using the NIST'11 and Wiley 10 mass spectral libraries [17-25]. Data are presented in tables 1 and 2.

Results and discussion

Table 1 - The data of chromatography-mass spectrometry of a hexane extract obtained from rose hips *Rosa canina L.*

Nº	Retention time	Name of compound	Gross formula	Quantity, %
1	26,89	Gentriacontan	C ₃₁ H ₆₄	3,00
2	28,46	Heptocazane	C ₂₇ H ₅₆	9,63
3	28,76	Bis 2 (ethyl hexyl) phthalate	C ₂₄ H ₃₈ O ₄	3,89
4	29,21	butyl tetradecyl ether hydrochloric acid	C ₁₈ H ₃₈ O ₃ S	3,75
5	29,93	Pentacosin	C ₂₅ H ₅₂	21,87
6	30,04	1-heptacosanol	C ₂₇ H ₅₆ O	14,69
7	30,63	Nonhexacontanoic acid	C ₃₈ H ₇₀ O ₂	1,99
8	31,32	Octadecan	C ₁₈ H ₃₈	19,16
9	31,48	Methyl 2-oxo octadecanoate	C ₃₉ H ₇₀ O ₃	3,10
10	31,64	Methyl -5.9.21-octa-cosatrienoate	C ₂₉ H ₅₂ O ₂	3,82
11	31,82	[4- (2-Methylpentanoylsulfamoyl) phenyl] amide 2-methylpentanoic acid	C ₁₈ H ₂₀ O ₄ S	3,00
12	32,97	Octadecanoic acid 2-oxo-methyl ester	C ₃₉ H ₇₀ O ₃	9,63

12 substances were identified in the hexane extract of the rosehip *Rosa canina L.*, of which Heptocazane, Pentacosin, Octadecane, belonging to the hydrocarbon class, have a high percentage, higher alcohols 1-Heptacosanol, derivatives of higher carboxylic acids (Methyl 2-oxo octadecanoate), Methyl-5,9,21-octa-cosatrienoate, 2-oxo-methyl ether of octadecanoic acid) and higher carboxylic acids Neagexacontanoic acid. In addition, Bis 2 (ethyl hexyl) phthalate (a phthalic acid derivative) and butyl tetradecyl ether of hydrochloric acid were identified in the extract.

Table 2 - Chromatography-mass spectrometry data of a chloroform extract obtained from rose hips of *Rosa canina L.*

№	Retention time	Name of compound	Gross formula	Quantity, %
1	28,47	2-methyl okatacosan	C ₂₉ H ₆₀	8,14
2	29,22	Butyl undecyl sulfuric acid ester	C ₁₅ H ₃₂ O ₃ S	3,23
3	29,93	Octadecan	C ₁₈ H ₃₈	21,23
4	30,04	Eicosyl trifluoroacetate	C ₂₄ H ₄₁ F ₃ O ₂	14,27
5	31,24	Naphthalene, decahydro-1,4a-dimethyl-7-(1-methylethyl)-, [1S-(1 α , 4 $\alpha\alpha$, 7 α , 8 $\alpha\beta$)] glucopyranosyloxy) -1-butenyl] -4	C ₁₅ H ₂₈	1,36
6	31,31	Octacosan	C ₁₈ H ₅₈	25,51
7	31,37	4-[3-(β -d-hydroxy-3,5,5-trimethyl-2-cyclohexen-1-one,	C ₂₇ H ₃₈ O ₁₂	1,54
8	31,82	3-benzyl-2,4-pentanedione	C ₁₂ H ₁₄ O ₁₂	4,46
9	32,97	1 eicosen	C ₂₀ H ₄₀	5,51
10	33,62	ethoxycarbonyl-3-methyl-4-azafluorenone, 2-fluorenilim	C ₂₉ H ₂₂ N ₂ O ₂	6,34

10 compounds were identified in the chloroform extract of rosehip *Rosa canina L.*

The dominant hydrocarbons are 2 methyl okatacosan, octadecane, octacosan and naphthalene, decahydro-1,4a-dimethyl-7-(1-methylethyl)-, [1S-(1 α , 4 $\alpha\alpha$, 7 α , 8 $\alpha\beta$)] glucopyranosyloxy) -1-butene] -4.

Separately, it should be noted that eicosyl trifluoroacetate belongs to the class of organohalogen compounds, which are relatively rare in plant objects. In addition, a homolog of butyl tetradecyl ether of hydrochloric acid was identified. Butyl undecyl ether of sulfuric acid was also detected in the hexane extract. Ethoxycarbonyl-3-methyl-4-azafluorenone, 2-fluorenilimime, which is a natural dye, was also identified in the chloroform extract.

Conclusion

For the first time, a comparative analysis of lipophilic substances contained in the fruits of the plant *Rosa canina L.*, harvested in the Almaty region, was studied and carried out.

The extracts mainly include hydrocarbons, derivatives of higher carboxylic acids, higher alcohols and some other substances, in particular butyl tetradecyl ether of hydrochloric acid, [4-(2-methylpentanoylsulfamoyl) phenyl] amide of 2-methylpentanoic acid in hexane extract and, butyl undecyl ether sulfuric acid, eicosyl trifluoroacetate and ethoxycarbonyl-3-methyl-4-azafluorenone, 2-fluorenilim in the chloroform extract.

Further work will be continued in order to determine the potential biological activity of the obtained extracts.

Е.С. Ихсанов, Г.Е. Тасмагамбетова, Ю.А. Литвиненко, Г.А. Сейтимова, Г.Ш. Бурашева

Әл-Фараби атындағы Қазак ұлттық университеті,
Физикалық-химиялық зерттеулер және талдау әдістері орталығы
Қазақстан Республикасы, Алматы, Қазақстан

ROSA CANINA L. ӨСІМДІК ЖЕМІСТЕРІНІҢ ЛИПОФИЛЬДІ ФРАКЦИЯСЫНЫН ФИТОХИМИЯЛЫҚ ҚУРАМЫ

Аннотация. Итмұрын немесе раушан (*Rosa L.*), Қызылт тұсті (*Rosaceae*) отбасына жатады. Жалпы Роза деп аталағын көптеген мәдени нысандары бар. Қазіргі уақытта итмұрының 400-ден астам түрі белгілі. Әдетте бұл тік бұталар, сирек Лиана, кейде биік емес ағаш тәріздес немесе шөпті өсімдіктер. Итмұрын

Солтүстік жарты шарның қалыпты және субтропикалық аймактарында тарапланған, оны тропикалық белдеудің таулы аудандарында кездестіруге болады. Итмұрының кейбір түрлері солтүстіктегі полярлық шенберден оңтүстіктегі Эфиопияға дейін жетеді. Ал Америка құрлығында-Канададан Мексикага дейін барады. Итмұрын үшін қолайлы жағдай Жерорта теңізі аймагы. Итмұрын түрінің бірнеше көң тарапалу аймагы бар.

Қазақстанда есімдік тектес дәрілік шикізаттың елеулі ресурстары бар. Жабайы өсетін флораның ең басым өкілдерінің қатарына итмұрын – *Rosa L.* тегінің түрлері жатады. Республикада итмұрының 21 түрі өседі, оның ішінде Орталық Қазақстанда-5: *R. glabrifolia* – бас сүйекті итмұрын, *R. laxa Retz.* – борпылдақ итмұрын, *R. acicularis Lindl.* – (ине итмұрын), *R. majalis Herrm.* (*R. cinnamomea L.*) – май итмұрын (сиыр итмұрын) и *R. pimpinellifolia L.* (*R. spinosissima L.*) – бедренцелистік итмұрын.

Итмұрын жемістері биологиялық белсененді заттардың жоғары болуымен ерекшеленеді және медицинада да, кулинарияда да кеңінен қолданылады.

Бұл жұмыста 2018 жылдың күздінде дайындалған итмұрын, *Rosa canina L.* жемістеріне салыстырмалы талдау жүргізілді.

Өсімдік шикізаты Қазақстан Республикасының оңтүстік өңірінде (Алматы облысында) жиналған. Ұсақталған ауа-құрғақ шикізат Сокслет аппаратындағы шикізат-реагент (1:10) қатынасында гексан және хлороформмен экстрагерленді. Алынған сығынды масс-селективті детекторы бар газды хроматографта талдаған қалың концентратка дейін жұмысашақ жағдайларда шоғырланылды.

Талдаулар 6890N/5973C (Agilent, АҚШ) масс-спектрометриялық детекторы бар газды хроматографта жүргізілді. Хроматограммаларда табылған шындарды сәйкестендіруді NIST'11 және Wiley 10 масс-спектрлері кітапханаларының көмегімен жүргізді.

Мақалада алғаш рет Алматы облысында дайындалған *Rosa canina L.* өсімдігінің жемістеріндегі липофильді заттарға салыстырмалы талдау жүргізілді және зерттелді.

Сығындылардың құрамына негізінен көмірсүтектер, жоғары карбон қышқылдарының туындылары, жоғары спирттер және басқа да заттар табылды, оның ішінде тұз қышқылының тетрадецил эфирі бутилі, [4- (2-метилпентаноилсульфамоил) фенил] гексан сығындысындағы 2-метилпентан қышқылы амиді және күкірт қышқылының бутил ундецил эфирі, Эйкозил трифторацетат және этоксикарбонил-3-метил-4 – азафлуоренон.

Одан кейінгі жұмыстар алынған сығындылардың әлеуетті биологиялық белсенділігін анықтау мақсатында жалғасатын болады.

Түйін сөздер: *Rosa canina L.*, ЖКФ-экстракция, никотин, GC-MS, химиялық құрамы.

Е.С. Ихсанов, Г.Е. Тасмагамбетова, Ю.А. Литвиненко, Г.А. Сейтимова, Г.Ш. Бурашева

Казахский национальный университет имени аль-Фараби,
Центр физико-химических методов исследования и анализа
Республика Казахстан, Алматы, Казахстан

ФИТОХИМИЧЕСКИЙ СОСТАВ ЛИПОФИЛЬНОЙ ФРАКЦИИ ПЛОДОВ РАСТЕНИЯ *ROSA CANINA L.* РОДА ШИПОВНИК (*ROSA*)

Аннотация. Род шиповник или роза (*Rosa L.*) относится к семейству розоцветных (Rosaceae). Имеет множество культурных форм, разводимых под общим названием Роза. На данный момент известны более 400 видов шиповника. Обычно это прямостоящие кустарники, реже лианы, иногда невысокие древовидной формы или почти травянистые растения. Шиповник распространён в умеренных и субтропических зонах Северного полушария, нередко его можно встретить в горных районах тропического пояса. Некоторые виды шиповника распространены от полярного круга на севере до Эфиопии на юге. На американском континенте - от Канады до Мексики. Благоприятнейшие условия для шиповника находятся в области Средиземноморья. Несколько видов рода шиповник имеют обширный ареал распространения .

Казахстан обладает значительными ресурсами лекарственного сырья растительного происхождения. К числу наиболее приоритетных представителей дикорастущей флоры относятся виды рода шиповник – *Rosa L.* как ценнейшего лекарственного растения. Всего в республике произрастает 21 вид шиповника, в том числе в Центральном Казахстане – 5: *R. glabrifolia* – ш. гололистый, *R. laxa Retz.* – ш. рыхлый, *R. acicularis Lindl.* – (ш. иглистый), *R. majalis Herrm.* (*R. cinnamomea L.*) – ш. майский (ш. коричный) и *R. pimpinellifolia L.* (*R. spinosissima L.*) – ш. бедренцелистый. Казахстанские виды рода *Rosa L.*

Плоды шиповника отличаются высоким содержанием биологически активных веществ и широко применяются как медицине, так и кулинарии.

В данной работе проводится сравнительный анализ плодов шиповника *Rosa canina L.*, заготовленного осенью 2018 года.

Растительное сырье собрано в южном регионе Республики Казахстан (Алматинская область). Измельченное воздушно-сухое сырье экстрагировали гексаном и хлороформом в соотношение сырье-реагент (1:10) в аппарате Сокслета. Полученный экстракт концентрировали в мягких условиях до густого концентрата, который анализировали на газовом хроматографе с масс-селективным детектором.

Анализы проводили на газовом хроматографе с масс-спектрометрическим детектором 6890N/5973C (Agilent, США). Идентификацию пиков, обнаруженных на хроматограммах, проводили при помощи библиотек масс-спектров NIST'11 и Wiley 10.

В статье впервые изучен и проведён сравнительный анализ липофильных веществ, содержащихся в плодах растения *Rosa canina L.*, заготовленного в Алматинской области.

В состав экстрактов входят преимущественно углеводороды, производные высших карбоновых кислот, высшие спирты и некоторые другие вещества, в частности, бутил тетрадециловый эфир соляной кислоты, [4-(2-метилпентаноилсульфамоил) фенил] амид 2-метилпентановой кислоты в гексановом экстракте и бутил ундесиловый эфир серной кислоты, Эйкоэзил трифторацетат и этоксикарбонил-3-метил-4-азафлуоренон, 2-флуоренилимим в хлороформенном экстракте.

Дальнейшие работы будут продолжены с целью определения потенциальной биологической активности полученных экстрактов.

Ключевые слова: *Rosa canina L.*, СКФ-экстракция, никотин, GC-MS, химический состав.

Information about authors:

Ikhsanov Yerbol Saginovich - PhD, Senior lecturer Department of Chemistry and Chemical Technology of the Al-Farabi Kazakh National University, Kazakh National University, e-mail: erbol.ih@gmail.com, <https://orcid.org/0000-0003-4640-9584>;

Litvinenko Yuliya Alekseevna - candidate of chemical sciences, Senior lecturer, Department of Chemistry and Chemical, Technology of the Al-Farabi Kazakh National University, e-mail: yuliya_litvinenk@mail.ru, <https://orcid.org/0000-0002-6387-187X>;

Seitimova Gulnaz Absattarovna - Ph.D. Senior lecturer at the Faculty Department of Chemistry and Chemical Technology of the Al-Farabi Kazakh National University, e-mail: sitigulnaz@mail.ru, <https://orcid.org/0000-0002-5157-1255>;

Burasheva Gauhar Shahmanovna - Doctor of Science, professor, Department of Chemistry and Chemical Technology of the Al-Farabi Kazakh National University, e-mail: gauharbur@mail.ru, <https://orcid.org/0000-0003-2935-3531>;

Tasmagambetova Gulzhaina Erzhanovna - Graduate student of the Faculty of Chemistry and Chemical Technology, e-mail: guljaina_98.14@mail.ru, <https://orcid.org/0000-0003-1892-2268>

REFERENCES

- [1] Nadpal J. D., Lesjak M. M., Šibul F. S., Anačkov G. T., Četojević-Simin D. D., Mimica-Dukić N. M., Beara I. N. (2016). Comparative study of biological activities and phytochemical composition of two rose hips and their preserves: *Rosa canina L.* and *Rosa arvensis Huds.* *Food Chemistry*, 192, 907–914. doi:10.1016/j.foodchem.2015.07.089
- [2] Ouerghemmi S., Sebei H., Siracusa L., Ruberto G., Saija A., Cimino F., & Cristani M. (2016). Comparative study of phenolic composition and antioxidant activity of leaf extracts from three wild *Rosa* species grown in different Tunisia regions: *Rosa canina L.*, *Rosa moschata Herrm.* and *Rosa sempervirens L.* *Industrial Crops and Products*, 94, 167–177. doi:10.1016/j.indcrop.2016.08.019
- [3] Ognyanov M., Remoroza C., Schols H. A., Georgiev Y., Kratchanova M., & Kratchanov C. (2016). Isolation and structure elucidation of pectic polysaccharide from rose hip fruits (*Rosa canina L.*). *Carbohydrate Polymers*, 151, 803–811. doi:10.1016/j.carbpol.2016.06.031
- [4] Winther K., Warholm L., Campbell-Tofte J., Marstrand K. (2018). Effect of *Rosa canina L.* (Rose-hip) on cold during winter season in a middle-class population: A randomized, double-blinded, placebo-controlled trial. *Journal of Herbal Medicine*. doi:10.1016/j.hermed.2018.04.003
- [5] Jakovljevic M., Moslavac T., Bilic M., Aladic K., Bakula F. (2018). Supercritical CO₂ extraction of oil from rose hips (*Rosa canina L.*) and cornelian cherry (*Cornus mas L.*) seeds. *Croatian Journal of Food Science and Technology*, 10(2), 197–205. doi:10.17508/cjfst.2018.10.2.08
- [6] Saricaoglu F. T., Atalar I., Yilmaz V. A., Odabas H. I., & Gul O. (2019). Application of multi pass high pressure homogenization to improve stability, physical and bioactive properties of rosehip (*Rosa canina L.*) nectar. *Food Chemistry*. doi:10.1016/j.foodchem.2019.01.002
- [7] Azmir J., Zaidul I.S.M., Rahman M.M., Sharif K.M., Mohamed A., Sahena F., Jahurul M.H.A., Ghafoor K., Norulaini N.A.N., Omar A.K.M. (2013). Techniques for extraction of bioactive compounds from plant materials: a review. *J. Food Eng*, vol. 117, no. 4, 426–436, sci-hub.tw/10.1016/j.jfoodeng.2013.01.014
- [8] Kou Y., Yuan C., Zhao Q., Liu G., Nie J., Ma Z., Zhao L. (2016). Thidiazuron Triggers Morphogenesis in *Rosa canina L.* Protocorm-Like Bodies by Changing Incipient Cell Fate. *Frontiers in Plant Science*, 7. doi:10.3389/fpls.2016.00557
- [9] Ikhsanov Y.S., Sultanova N.A., Abilov Z.A., Choudhary M.I. (2018). Investigation of components of hexane extract from *Tamarix hispida* by method of gas chromatography. *Izvestiya vysshikh uchebnykh zavedenii khimiya khimicheskaya tekhnologiya*, vol.61, no6, 83-87, doi: <https://doi.org/10.6060/tctc.20186106.5706>
- [10] Gruenwald J., Uebelhack R., Moré M. I. (2019). *Rosa canina* - Rose Hip Pharmacological Ingredients and Molecular Mechanics Counteracting Osteoarthritis – a Systematic Review. *Phytomedicine*, 152958. doi:10.1016/j.phymed.2019.152958

- [11] Jafarirad S., Mehrabi M., Divband B., Kosari-Nasab M. (2016). *Biofabrication of zinc oxide nanoparticles using fruit extract of Rosa canina and their toxic potential against bacteria: A mechanistic approach.* Materials Science and Engineering: C, 59, 296–302. doi:10.1016/j.msec.2015.09.089
- [12] Jiménez S., Gascón S., Luquin A., Laguna M., Ancin-Azpilicueta C., Rodríguez-Yoldi M. J. (2016). *Rosa canina Extracts Have Antiproliferative and Antioxidant Effects on Caco-2 Human Colon Cancer.* PLOS ONE, 11(7), e0159136. doi:10.1371/journal.pone.0159136
- [13] Ikhsanov Y.S., Nauryzbaev M., Musabekova A., Alimzhanova M., Burashev E.(2019). Study of nicotiana tabacum l extraction, by methods of liquid and supercritical fluid extraction, journal-Journal of Applied Engineering Science, Vol.17, no. 3 doi:10.5937/jaes17-21189
- [14] Taghizadeh M., Rashidi A. A., Taherian A. A., Vakili Z., Sajad Sajadian M., Ghardashi M. (2016). Antidiabetic and Antihyperlipidemic Effects of Ethanol Extract of Rosa canina L. fruit on Diabetic Rats. Journal of Evidence-Based Complementary & Alternative Medicine, 21(4), NP25–NP30. doi:10.1177/2156587215612626
- [15] Erfani M., Tabatabaei Z. G., Sadigh-Eteghad S., Farokhi-Sisakht F., Farajdokht F., Mahmoudi J., Nasrolahi A. (2019). *Rosa canina L. methanolic extract prevents heat stress-induced memory dysfunction in rat.* Experimental Physiology. doi:10.1113/ep087535
- [16] Kiralan M., Yildirim G. (2019). *Rosehip (Rosa canina L.) Oil. Fruit Oils: Chemistry and Functionality,* 803–814. doi:10.1007/978-3-030-12473-1_43
- [17] Kilinc K., Demir S., Turan I., Mentese A., Orem A., Sonmez M., Aliyazicioglu Y. (2019). *Rosa canina Extract has Antiproliferative and Proapoptotic Effects on Human Lung and Prostate Cancer Cells.* Nutrition and Cancer, 1–10. doi:10.1080/01635581.2019.1625936
- [18] Valcheva-Kuzmanova, S., Denev, P., Eftimov, M., Georgieva, A., Kuzmanova, V., Kuzmanov, A., ... Tzaneva, M. (2019). *Protective effects of Aronia melanocarpa juices either alone or combined with extracts from Rosa canina or Alchemilla vulgaris in a rat model of indomethacin-induced gastric ulcers.* Food and Chemical Toxicology, 132, 110739. doi:10.1016/j.fct.2019.110739
- [19] Hu, X.-Y., Corp, N., Quicke, J., Lai, L., Blondel, C., Stuart, B., ... Moore, M. (2018). *Rosa canina fruit (rosehip) for osteoarthritis: a cochrane review.* Osteoarthritis and Cartilage, 26, S344. doi:10.1016/j.joca.2018.02.685
- [20] Chemat, F., Vian, M. A., & Cravotto, G. (2012). Green Extraction of Natural Products: Concept and Principles. International Journal of Molecular Sciences, vol.13, no.7, 8615–8627. doi:10.3390/ijms13078615
- [21] Farajpour, R., Sadigh-Eteghad, S., Ahmadian, N., Farzipour, M., Mahmoudi, J., & Majdi, A. (2017). *Chronic Administration of Rosa canina Hydro-Alcoholic Extract Attenuates Depressive-Like Behavior and Recognition Memory Impairment in Diabetic Mice: A Possible Role of Oxidative Stress.* Medical Principles and Practice, 26(3), 245–250. doi:10.1159/000464364
- [22] Al-Yafeai A., Malarski A., Böhm V. (2018). *Characterization of carotenoids and vitamin E in R. rugosa and R. canina : Comparative analysis.* Food Chemistry, 242, 435–442. doi:10.1016/j.foodchem.2017.09.070
- [23] Nowrouzi F., Azadbakht M., Kalehoei E., Modarresi M. (2019). *Protective Effect of Rosa Canina Extract Against Doxorubicin-induced Testicular Toxicity in Mice.* Brazilian Archives of Biology and Technology, 62. doi:10.1590/1678-4324-2019180017
- [24] Fattah A., Niyazi F., Shahbazi B., Farzaei M. H., Bahrami, G. (2016). *Antidiabetic Mechanisms of Rosa canina Fruits.* Journal of Evidence-Based Complementary & Alternative Medicine, 22(1), 127–133. doi:10.1177/2156587216655263
- [25] Myrzakhanova M. N., Kaa S. A. New trends of medical media in training and practice // NEWS of the National academy of sciences of the Republic of Kazakhstan. Series of biological and medical. 2018. Vol. 5, N 329. P. 67-70. (in Eng). <https://doi.org/10.32014/2018.2518-1629.9>

Publication Ethics and Publication Malpractice in the journals of the National Academy of Sciences of the Republic of Kazakhstan

For information on Ethics in publishing and Ethical guidelines for journal publication see <http://www.elsevier.com/publishingethics> and <http://www.elsevier.com/journal-authors/ethics>.

Submission of an article to the National Academy of Sciences of the Republic of Kazakhstan implies that the described work has not been published previously (except in the form of an abstract or as part of a published lecture or academic thesis or as an electronic preprint, see <http://www.elsevier.com/postingpolicy>), that it is not under consideration for publication elsewhere, that its publication is approved by all authors and tacitly or explicitly by the responsible authorities where the work was carried out, and that, if accepted, it will not be published elsewhere in the same form, in English or in any other language, including electronically without the written consent of the copyright-holder. In particular, translations into English of papers already published in another language are not accepted.

No other forms of scientific misconduct are allowed, such as plagiarism, falsification, fraudulent data, incorrect interpretation of other works, incorrect citations, etc. The National Academy of Sciences of the Republic of Kazakhstan follows the Code of Conduct of the Committee on Publication Ethics (COPE), and follows the COPE Flowcharts for Resolving Cases of Suspected Misconduct (http://publicationethics.org/files/u2/New_Code.pdf). To verify originality, your article may be checked by the Cross Check originality detection service <http://www.elsevier.com/editors/plagdetect>.

The authors are obliged to participate in peer review process and be ready to provide corrections, clarifications, retractions and apologies when needed. All authors of a paper should have significantly contributed to the research.

The reviewers should provide objective judgments and should point out relevant published works which are not yet cited. Reviewed articles should be treated confidentially. The reviewers will be chosen in such a way that there is no conflict of interests with respect to the research, the authors and/or the research funders.

The editors have complete responsibility and authority to reject or accept a paper, and they will only accept a paper when reasonably certain. They will preserve anonymity of reviewers and promote publication of corrections, clarifications, retractions and apologies when needed. The acceptance of a paper automatically implies the copyright transfer to the National Academy of Sciences of the Republic of Kazakhstan.

The Editorial Board of the National Academy of Sciences of the Republic of Kazakhstan will monitor and safeguard publishing ethics.

Правила оформления статьи для публикации
в журнале смотреть на сайте:

www:nauka-nanrk.kz

<http://chemistry-technology.kz/index.php/en/arhiv>

ISSN 2518-1491 (Online), ISSN 2224-5286 (Print)

Редакторы: *М. С. Ахметова, Г. Б. Халидуллаева, Д. С. Аленов*
Верстка на компьютере *А.М. Кульгинбаевой*

Подписано в печать 29.03.2020.
Формат 60x881/8. Бумага офсетная. Печать – ризограф.
10 п.л. Тираж 300. Заказ 2.

Национальная академия наук РК
050010, Алматы, ул. Шевченко, 28, т. 272-13-18, 272-13-19